

Concentrations of mercury in milligrams per kilogram (mg/kg) wet weight in walleye fillet tissue from the upper and lower reaches of Lake Roosevelt from the 1994 and 1998 USGS studies.

[cm, centimeters; mg/kg, milligrams per kilogram; ww, wet weight; Upper, between Northport and Kettle Falls; Lower, between Spokane River and Grand Coulee Dam]

1994 USGS study			1998 USGS study		
Reach	Length (cm)	Mercury (mg/kg, ww)	Reach	Length (cm)	Mercury (mg/kg, ww)
Upper	36.2	0.14	Upper	36.0	0.12
Upper	34.7	0.16	Upper	35.0	0.10
Upper	37.4	0.20	Upper	33.5	0.11
Upper	34.3	0.15	Upper	36.0	0.10
Upper	34.3	0.22	Upper	36.0	0.11
Upper	40.4	0.30	Upper	33.5	0.12
Upper	38.1	0.16	Upper	35.0	0.12
Upper	36.5	0.18	Upper	34.5	0.12
Lower	34.1	0.31	Lower	36.0	0.13
Lower	35.6	0.21	Lower	34.0	0.16
Lower	34.5	0.24	Lower	36.5	0.12
Lower	36.2	0.26	Lower	37.5	0.15
Lower	35.8	0.18	Lower	40.5	0.19
Lower	35.1	0.28	Lower	39.5	0.22
Lower	34.6	0.27	Lower	41.0	0.18
Lower	36.6	0.30	Lower	35.5	0.10